SCORING MANUAL FOR THE QUALITY OF LIFE IN EPILEPSY INVENTORY-89 (QOLIE-89)

CONTENT OF QOLIE-89 (Table 1)

QOLIE-89 contains I7 multi-item scales that tap the following health concepts: overall quality of life (2 items), emotional well-being (5 items), role limitations due to emotional problems (5 items), social support (4 items), social isolation (2 items), energy/fatigue (4 items), seizure worry (5 items), medication effects (3 items), health discouragement (2 items), work/driving/social function (11 items), attention/concentration (9 items), language (5 items), memory (6 items), physical function (10 items), pain (2 items), role limitations due to physical problems (5 items), and health perceptions (6 items). A QOLIE-89 overall score is obtained using a weighted average of the multi-item scale scores. QOLIE-89 also includes one item on change in health over the preceding year and two items added after field testing: one on overall health and one on satisfaction with sexual relations.

The generic core of QOLIE-89 is the RAND 36-Item Health Survey 1.0, also known as the SF-36 (Ware and Sherbourne, 1992; Hays, Sherbourne, and Mazel, 1993). Items in this 36-item measure were adapted from longer instruments completed by patients participating in the Medical Outcomes Study (MOS), an observational study of variations in physician practice styles and patient outcomes in different systems of health care delivery (Stewart, Sherbourne, Hays, et al, 1992).

In addition to the generic core, 13 items from longer MOS instruments and 5 items originally developed for the Epilepsy Surgery Inventory-55 (Vickrey, Hays, Graber, et al, 1992) were incorporated into QOLIE-89. The two-item overall quality-of-life scale consists of one Dartmouth COOP Chart (Nelson, Landgraf, Hays, et al, 1990) and one item from a study on patient preferences (Hadorn and Hays, 1991). This latter item was itself adapted from the Faces Scale (Andrews and Withey, 1976). The single item on overall health was adapted from an existing visual analog scale (Brazier, Jones, and Kind, 1993).

The remaining 32 items were developed *de novo* by the QOLIE Development Group based on diverse clinical experience with patients and a review of the literature on patient concerns about health-related quality of life.

FIELD TESTING

Item selection for QOLIE-89 was based on analysis of data collected from a cohort of 304 adult men and women having simple partial, complex partial, grand mal, absence, and/or myoclonic seizures of mild to moderate severity. These patients were enrolled from 25 sites across the United States.

All subjects completed an initial 98-item QOLIE test battery; the majority of subjects completed this same battery again within 3 weeks of the first visit. A brief neuropsychological test battery, selected neurological exam features, a proxy's assessment of the subject's quality of life, and information about seizure occurrence, medications, demographic characteristics, and health care utilization were also obtained (Perrine, 1993).

Data from this study were analyzed, and three measures of quality of life were developed, differing in their number of items: QOLIE-89, QOLIE-31 (Vickrey, Perrine, Hays, et al., 1993), and QOLIE-10.

SCORING RULES

Seventeen Primary Scales. Precoded numeric values for responses on some QOLIE-89 items are in the direction such that a higher number reflects a more favorable health state. For example, a circled response of "10" for item 2 corresponds to best possible quality of life, while a circled response of "0" corresponds to worst possible quality of life. However, precoded numeric values for some other items are in the direction such that a *lower* number reflects a more favorable health state. For example, a circled response of "1" for item 49 corresponds to a more favorable quality of life, while a value of "5" on this item corresponds to a less favorable quality of life. As these examples also demonstrate, different items have different ranges of precoded numeric values.

To account for these differences, the scoring procedure for QOLIE-89 first converts the raw precoded numeric values of items to 0-100 point scores, with higher converted scores always reflecting better quality of life (Table 2). To perform this step, write in the converted score for each item in the column labeled "Subtotal" in Table 2. Next, sum the subtotal scores for each scale and write in these values in the places marked "Total." Finally, divide each "Total" by the number of items that the respondent answered within each scale to get the "Final Score." The possible range of each scale's final score is now from 0 to 100 points. Higher scores reflect better quality of life; lower ones, worse quality of life.

Note that Table 2 shows the divisors to be used only in situations where every item within a given scale has been answered. For example, if item 40 in the Seizure Worry scale was left blank and the other four items in the scale were answered, then the "Total" score for Seizure Worry would be divided by "4" (instead of "5") to obtain the "Final Score."

Overall Score. A QOLIE-89 overall score can be derived by weighting and summing QOLIE-89 scale scores (Table 3). QOLIE-89 scale weights were derived in the following way:

A factor analysis of the 17 QOLIE-89 scales was performed. A four-factor solution yielded unique placement of scales into domains of mental health, physical health, cognitive function, and epilepsy-specific areas, based on the magnitude of each scale's factor loadings. Factor scores, which weight each scale by its factor loading, were derived for each domain. These four factor scores were then averaged to produce a single summary score.

To derive QOLIE-89 scale weights, this summary score was regressed on the I7 QOLIE-89 multi-item scales. Standardized beta coefficients from this regression analysis were summed, and each beta coefficient was divided by the sum to derive the relative weight for each QOLIE-89 scale listed in Table 3.

The QOLIE-89 scale overall score is calculated by summing the product of each scale final score from Table 2 times its weight and summing all these subtotals (Table 3).

T-scores. T-scores can be determined for each of the 17 scale final scores (Table 2) and for the overall score (Table 3) using the key in Table 4. These T-scores represent linear transformations of the scores that produce a mean of 50 and standard deviation of 10 for the cohort of 304 adults with epilepsy. Thus, a person with a T-score of 50 has a score equal to that of the mean for the epilepsy cohort. T-scores were computed using the following formula:

T-score =
$$50 + \left[10 \left(\frac{\text{Observed final scale score } minus \text{ scale mean in Table 5}}{\text{Scale standard deviation in Table 5}} \right) \right]$$

Higher T-scores reflect a more favorable quality of life.

RELIABILITY AND DESCRIPTIVE STATISTICS

Table 5 presents reliability coefficients and descriptive statistics for the 17 QOLIE-89 scales derived from analyses of the 304 respondents in the QOLIE development study. Internal consistency reliabilities (Cronbach's alpha) range from r=0.78 to r=0.92, exceeding the 0.70 standard for group-level comparisons (Nunnally, 1978) for all scales. Test-retest reliabilities were calculated as Pearson product-moment correlations between patient responses at visit 1 and patient responses at visit 2 (up to 3 weeks later). At the second testing, patients were asked whether they had experienced any major life-and-death or health-related events since the first visit; data on those patients responding in the affirmative were excluded from test-retest analyses. The test-retest reliabilities for the 17 scales range from r=0.58 to r=0.86. All scales, except role limitations due to physical problems (r=0.58), role limitations due to emotional problems (r=0.67), medication effects (r=0.64), and pain (r=0.69), exceeded the r=0.70 standard for group comparisons. The 3 scales having the lowest test-retest reliabilities also had the 3 largest standard deviations among the 17 scales. The overall score had an internal consistency reliability of 0.97 and test-retest reliability of 0.88.

Means of the 17 self-report scales range from 54.3 for the memory scale to 85.3 for the physical function scale. None of the scales show floor or ceiling effects (scores clustering near the minimum or maximum). Examination of the means and associated standard deviations and ranges show sufficient breadth to assess a broad range of functioning and the potential to detect changes in quality of life.

TABLE 1
KEY TO SOURCES OF ITEMS IN QOLIE-89

Item Number	Source
2	Adapted from the Faces Scale (Andrews and Withey, 1976) by Hadorn and Hays (1991)
1,3,4-17,19,20, 23-35,43*47	From RAND 36-Item Health Survey 1.0 (aka SF-36) (Ware and Sherbourne, 1992; Hays, Sherbourne and Mazel, 1993)
18,21,22,40,50	From Epilepsy Surgery Inventory (ESI)-55 (Vickrey, Hays, Graber, et al, 1992)
37,38,39,41,48,64,73 81-83,86-88	From longer instruments in the Medical Outcomes Study (Stewart, Sherbourne, Hays, et al, 1992)
49	Dartmouth COOP Chart (Nelson, Landgraf, Hays, et al., 1990)
36,42,51-63,65-72, 74-80,84,85	Developed de novo by QOLIE Development Group
89*	Visual analog item adapted from existing measure (Brazier, Jones, and Kind, 1993)

^{*}The wording of items 43 and 89 in the QOLIE-89 Inventory has been modified slightly since field testing and publication of the QOLIE-31 Scoring Manual.

TABLE 2
QOLIE-89 SCORING FORM

Response (raw score)								Final Score,		
Scale/Item_Numbers	1	2	3	4	5	6	Subtotal	0-100 point scale		
Health Perceptions			_		•					
1.	100	75	(50)	25	0	_	-			
44.	0	25	(50)	75	100	_				
45 .	100	75	50	(25)	0	_				
46.	0	(25)	50	75	100					
47.	100	75	50	25)	0	_				
48.	0	25	50	75	100	_				
40.	U	25	50	(13)	100	TOTAL:	250 ÷	6 = 41.67		
Overall Quality of Life										
2.	(multip	y raw so	core by	(10)			40			
49.	100	75	50	25	0	_	.400			
40.	100	, 0	50		Ü	TOTAL:	<u>69</u> ÷	2 = 32.5		
Physical Function							•			
4.	ကြိမ်ဝ	50	100		_	_				
5 .	X	50	100	_	_	_				
6.	Ō		100	_						
7.	Ö	50 50	100	_	_	_				
8.	Ö	50	100							
9.		7		_						
	0	50 50 50	100		_	_				
10.	0	56	100	_	_	_				
11.	0	<u>650</u>	100	_			***************************************			
12.	0	50	100	_	_	_				
13.	0	50	100	-	_	— TOTAL		÷ 10= 55		
						TOTAL:	<u>550</u> -	+ 10= <u>55</u>		
Role Limitations-Physic	cal	\sim								
14.	0	(100	_	_			-			
15.	0	100			_	_				
16.		100	_	_	_	_				
17.		100	_	_						
18.	0	TOD	_	_		_	***************************************	4.4		
10.	U	100		_		TOTAL:	300	+ 5 = <u>40</u>		
Role Limitations-Emot	ional									
19.		100	_			_				
20.	70	100				_				
21.							······································			
		100			_	_	***************************************			
22.	9999	100			_	_				
23.	0	100			_	TOTAL:	<u>O</u> -	÷ 5 = <u></u>		
						IOIAL.	entral manifestation and the second	- 0 = <u>-</u>		
Pain	_									
24.	100	80	60	40	20	0	· <u></u>			
25.	100	75	50	25	0	_				
						TOTAL:	700	÷ 2 = <u>CO</u>		

TABLE 2 QOLIE-89 SCORING FORM (cont.)

		Response (raw score)						Final Score,	
Scale/Item Numbers	1	2	3	4	5	6	Subtotal	0-100 point scale	
Work/Driving/									
Social Function									
26 .	100	75	50	25	0	_			
36.	0	20	40	60	80	100			
43.	0	20	40	60	80	100	·		
65.	0	2 5	50	75	100				
66.	90	25	50	75	100	_			
67.	(25	50	75	100	_			
68.	0	25	(50)	75	100		· **********		
76.	100	75	(50)	25	0	_			
7 7.	100	75	50	25					
78.	100	75	50	25	(•		
85.	0	25	50	(75)	100	_	<u>-</u>	5	
						TOTAL:	24 S ÷	- 11= 22,28	
Energy/Fatigue									
27.	100	80	60	40	20	0			
31.	100	80	60	40	20	0	-		
33.	0	20	40	60	80	100			
35.	0	20	40	60	80	100			
55 .	U	20	40)	60	80	TOTAL:	1612 ÷	-4 = 40	
Emotional Well-Being								***************************************	
28.	0	(20)	40	00	00	100			
20. 29.	0 0	20	40	60	80	100			
29. 30.	100		40	60	80	100			
30. 32.		80	60	40	20	0			
32. 34.	0	20	40	60	80	100			
34.	100	80	60	40	20	0 TOTAL:	<u>40</u> +	- 5 = 2 2	
						IOIAL.	•	5 = 4	
Attention/Concentration	l			_					
37.	0	20	40	(60)	80	100			
38 .	0	20	40	60	80	100			
41.	0	20	40	60	80	100			
60.	0	20	40	60	80	100			
6 1.	0	20	40	60	80	100			
62.	0	20	40	60	80	100			
63.	0	20	40	60	80	100			
64.	0	20	40	60	80	100			
73.	0	25	50	75)	100			0 0	
						TOTAL:	795_ +	e 9 = <u>88.3</u>	
Health Discouragement									
39.	0	20	40	60	80	100			
42.	0	20	40	60	80	100	<		
7 2.	U	20	40	00	OU	TOTAL:	40	- 2 = <u>20</u>	
						IOIAL.	70 7	- 2 = <u>~~</u>	

TABLE 2 QOLIE-89 SCORING FORM (cont.)

	Response (raw score)					Final Score,		
Scale/Item Numbers	1	2	3	4	5	6	Subtotal	0-100 point scale
Seizure Worry								
40.	0	20	40	60	80	100		
69 .	0	33.3	66.7	100		_		
70.	0	(50)	100	_			-	
71.	0	33.3	66.7	100	_			
74.	100	33.3 75	50	25	0			5 = 7168
						TOTAL:	33.4 ÷	· 5 = <u>/ </u>
Memory				_				_
50.	0	33.3	66.7	100	_	_	. *************************************	
51.	0	20	40	60	80	100		
52.	0	20	40	60	80	100		
53.	0	20	40	60	80	100	· ///////	
<u>54</u> .	0	20	40	60	80	(100)		
75 .	100	75	50	25	0		(- (~~)	- 6 = <u>[O</u>
						IOIAL:	<u>600</u> ÷	6 = 00
Language	_					<u>a</u>		
55 .	0	20	40	60	80	(100)	·	
56.	0	20	40	60	80	1990		
57.	0	20	40	60	80	100	· 	
58. 59.	0 0	20	40	60	80	100		
35.	U	20	40	60	80	TOTAL:	TOD -	- 5 = 100
Madiantian Effects						TOTAL.		J =
Medication Effects	1	00.0	00.7	400				
72. 79.	100	33.3	66.7	100	<u></u>	_		
79. 80.	100	75 75	50 50	25 25	0	_		
00 .	100	75	50	25	<i>,</i> 0	TOTAL ·	25	+ 3 = <u>33</u> 3
Secial Support						.0., 12.		
Social Support 81.	0		50	75	400			
81. 82.	0 0	25	50 30	75 75	100 100	*****		
83.	0	25 25	50	75 7 <u>5</u>	100	_		
86.	100	75	50 50	25	0	-		
00 .	100	73	30	23	U		12C -	+ 4 = <u>31.2</u> 5
Coalel Isolatin-							1 65-	· <u></u>
Social Isolation	0	(20)	40	-00	00	400		
87.	0	20	40	60	80	100		
88.	U	20	40	60	80	100 TOTAL:	<u> </u>	+ 2 <u>-20</u>
SINGLE ITEMS:						IOIAL.	<u> </u>	F 2 =
Change in Health		0						
3.	100	(75)	50	25	0			7 <u>5</u>
Cavual Dalatians								
Sexual Relations	100	75	EO	05				
84.	100	75	50	25		_		
Overall Health								
89.	(no red	coding n	ecessar	y)				
		-						

Note: The total number of items in each scale is listed as the divisor for each subtotal. However, where all items in a scale are not answered, the divisor will be lower, as noted in the text for "Scoring Rules," page 4.

TABLE 3
FORMULA FOR CALCULATING QOLIE-89 OVERALL SCORE

QOLIE-89 Scale	Final Score (from Table 2)		Weight		Subtotal
Health Perceptions	41.67	×	.06	=	2.5 (a)
Overall Quality of Life	32,5	×	.06	=	2.0 (b)
Physical Function	55	×	.06	=	3.3 (c)
Role Limitations-Physical	40	×	.07	=	2.8 (d)
Role Limitations-Emotional	<u>O</u>	×	.05	=	<u></u> (e)
Pain	100	×	.07	=	2 (f)
Work/Driving/Social Function	22.28	×	.0è	=	(<u>.</u> 8 (g)
Energy/Fatigue	<u>40</u>	×	.05	=	2 (h)
Emotional Well-Being	2 <u>8</u>	×	.05	=	1 <u>4</u> (i)
Attention/Concentration	88.3	×	.08	=	7 <u>, (j)</u>
Health Discouragement	20	×	.07	=	1. <u>4</u> (k)
Seizure Worry	71.68	×	.06	=	4 <u>3</u> (1)
Memory	100	×	.07	=	<u> </u>
Language	100	×	.06	=	<u>(n)</u>
Medication Effects	8.33	×	.05	=	0.4 (0)
Social Support	31.25	×	.02	=	<u>06</u> (p)
Social Isolation	20	×	.04	=	0 _ 3 _ (q)
OVERALL SCORE: Sum subtotal:	s (a) through (q)			=	41.9

TABLE 4 QOLIE-89 PROFILE SHEET

Т	Health Perceptions	Overall Quality of Life	Physical Function	Role-Physical	Role-Emotional	Pain	T
73							73
72 71							72 71
70					<u> </u>		70
69							69
68		100					68
67	100	98			ļ		67
66 65	100	95					66 65
64	96	93					64
63							63
62	92	90					62
61 60	88	88 85	······································			100	61
59	00	83		. 100	100	100	60 59
58	83			1	100	·	58
57		80	100				57
56		78				90	56
55 54	79	75	95	90		88	55
53	75	75 73		80	80	 	54 53
52		70	90			80	52
51	71					78	51
50		68	85			75	50
49 48	67	65 63				30	49
47	63	03	80	60	60	70 68	48 47
46		60				65	46
45	58	58	75			63	45
44						60	44
43	54	55 53	70	40	45	58	43
41	50	50	70	40	40	55 53	42 41
40		48	65			50	40
39	46					48	39
38		45				45	38
37	42	43	60			43	37
36 35	38	40	55	20	20	40	36 35
34		38	93			38 35	34
33						33	33
32	33	35	50			30	32
31		33					31
29	29	30 28	45	0	0	25	30
28	25					23 20	29 28
27		25	40				27
26	21	23					26
25			35			13	25
24	17	20 18				10	24
22	13	15	30			1	23 22
21	<u>'</u>	13					21
20			25			0	20
19	8	10					19
18	ļ						18
17 16	4	5	20	<u> </u>			17
15	0	5	15	- 		-	16 15
14	Ž.		.,			-	14
13		0					13
12			10				12
11			<u> </u>				11
10 9			5			<u> </u>	10
8							9
7			0				7
6		· · · · · · · · · · · · · · · · · · ·			+ 		6

TABLE 4 QOLIE-89 PROFILE SHEET (cont.)

7 73	Social Function	Energy/Fatigue	Emotional Well-Being	Attention/Concentration	Health Discouragement	Seizure Worry	1 7
72							7
71		100					7
70		100					7
9		95					7
86		33					6
57			100				6
6		90	100				6
35		30				100	6
34	98,100	85	96	100		96	6
33	00,100	- 65		98		95	6
52	95	80	92	96-97		91-93	6
31	91,93	80		94-95		88,90	6
60	89		88	92-93	100	86-87	6
9		75		90-91		83-85	6
8	86	73	84	88-89		81-82	5
7	82.84	70		86-87		78-80	5
6	80	70	80	84-85	90	76-77	5
5		CE		82-83		73-75	5
4	75,77	65	76	80-81		70,72	5
3	73	·····		78	80	68-69	5
2	13			76		65,67	5
1	70	60	72			63-34	5
0	66,68	<i></i>		73		60-61	5
9		55	68	69,71	70	59	5
8	64			67		55-57	4
7	50.04		64			52-53	4
	59,61	50		64		50-51	4
6	57		60	62	60	47-48	4
5	55	45		60		46	4:
4	52		56	58		42,44	4
3	50	40		56	50	40	
2	48		52	53		38	4
1				51			4:
0	43,45	35	48	49		34,36	4
9	41			47	40	32-33	41
8	39	30	44			29-30	35
7	36			44		27-28	38
6	34	25	40	40,42	30	24-25	3
5	32			38	30	21,23	3€
4	30		36	36		19-20	35
3	27	20				16-17	34
2	25		32	33		15	3:
1	24	15		31	20	11-13	32
D	20,22		28	29			3.
9	18	10		27		7-8	3(
В	16		24			4	29
7	15			24	10		28
5	11,13	5	20	22		0	27
5	9		20	20			26
4	7	0		18	0		25
3	5		16	16			24
2	2		16				23
-	ō		40	11,13			22
-			12	9			21
,				7			20
3			8				19
,				4			18
-			4	2			17
				0 1			16
5		<u> </u>	0				15
-							
} [· · · · · · · · · · · · · · · · · · ·						14
			-				13
							12
	<u>i</u>						11
)		ļ		•			
							10
)							10 9
)							

Note: If the final score is not on this table, either interpolate it or calculate, it directly using the formula on page 4.

TABLE 4 QOLIE-89 PROFILE SHEET (cont.)

73	Memory	Language	Medication Effects	Social Support	Social Isolation	Overall Score	Ť
72			<u> </u>				73
71					<u> </u>	<u> </u>	72
70				<u></u>		100	71
69	100					99	70
68	97					97-98	69
67	96					96	68
6 6	92-94					94-95	67
65	90		100			93	66
64	87-89				·	91-92 89-90	65
63	85-86			· · · · · · · · · · · · · · · · · · ·		88	63
62	83-84	100	92	100		86-87	62
61	80-82		89			85	61
60	78-79	96				83-84	60
59	75-77	· · · · · · · · · · · · · · · · · · ·	83	94	100	82	59
58	73-74	92	81	,		80-81	58
57	71-72		78	88		79	57
56	68-70	88	75			77-78	56
5 5	66-67		72		90	75-76	55
54 53	63-64	84	67.69	81		74	54
52	61-62	80	64			72-73	53
51	58-60 56-57	7.0	61		<u> </u>	71	52
50	56-57 54	76	58	75	80	69-70	51
49	51-53	70	56			68	50
48	49-50	72	53			66-67	49
47	46-48	60	50	69		65	48
46	44	68	47		70	63-64	47
45	42-43	64	42.44	63		61-62	46
44	39-41		39			60	45
43	37-38	60	36 33	56		58-59	44
42	34,36		31	30	60	57	43
41	32-33	56	28			55-56	42
40	29-31		25	50		54	41
39	27-28	52	22			52-53	40
38	26		19	44	50	51	39
37	22-24	48	17	77		49-50	38
36	20-21					47-48	37
35	17-19	44	11	38	40	46	36
34	16	40	8		70	44-45	35
33	13-14					43 41-42	34
32	10-12	36	0	31		40	33
31	8-9				30	38-39	31
30	6-7	32				37	30
29	3			25		35-36	29
28		28				33-34	28
27	0			19	20	32	27
26		24				30-31	26
25						29	25
24		20		13		27-28	24
23					10	26	23
22		16				24-25	22
21				6		23	21
20		12				21-22	20
19					0	19-20	19
18		8		0		18	18
17						16-17	17
16 15		4				15	16
15						13-14	15
13		0				12	14
13						10-11	13
						9	12
11						7-8	11
10 9						5-6	10
8						4	9
0						2.3	8
7	l l						
7						0	7

Note: If the final score is not on this table, either interpolate it or calculate it directly using the formula on page 4.

TABLE 5
RELIABILITY, CENTRAL TENDENCY, AND VARIABILITY OF QOLIE-89 SCALES*

		Re	eliability			
Scale	Number of Items	Alpha	Test-retest ^b	Mean (0-100 range)	Standard Deviation	Observed Range
Health Perceptions	6	0.78	0.84	68.26	19.61	21-100
Overall Quality of Life	2	0.79	0.84	67.17	18.38	5-100
Physical Function	10	0.89	0.75	85.27	19.82	6-100
Role Limitations-Physical	5	0.81	0.58	67.81	34.53	0-100
Role Limitations-Emotional	5	0.81	0.67	69.29	34.54	0-100
Pain	2	0.87	0.69	75.56	24.80	0-100
Work/Driving/Social Function	11	0.86	0.86	66.91	22.94	4-100
Energy/Fatigue	4	0.84	0.75	55.30	21.10	0-100
Emotional Well-Being	5	0.83	0.77	67.20	19.28	16-100
Attention/Concentration	9	0.92	0.86	69.98	20.70	11-100
Health Discouragement	2	0.82	0.73	69.87	27.74	0-100
Seizure Worry	5	0.79	0.84	58.29	25.76	0-100
Memory	6	0.88	0.82	54.34	24.15	0-100
Language	5	0.88	0.72	74.57	20.99	4-100
Medication Effects	3	0.78	0.64	55.34	30.52	0-100
Social Support	4	0.84	0.78	72.47	22.89	0-100
Social Isolation	2	0.88	0.73	76.78	25.04	0-100
Overall Score	86	0.97°	0.88	67.90	15.55	26-95

aN ranged from 298 to 304 patients with mild or moderate epilepsy for all data except test-retest reliability.

^bN ranged from 229 to 232 in the subset of epilepsy patients who were clinically stable and whose test-retest interval ranged from 1 to 21 days.

cEstimated using Mosier's (1943) formula.

REFERENCES

Andrews FM, Withey SB. Social Indicators of Well-Being: Americans' Perception of Life Quality. New York, NY: Plenum Press, 1976.

Brazier J, Jones J, Kind P. Testing the validity of the Euroquol and comparing it with the SF-36 health survey questionnaire. *Quality of Life Research.* 1993;2:169-180.

Hadorn D. Hays RD. Multitrait-multimethod analysis of health-related quality of life preferences. *Med Care.* 1991;29:829-840.

Hays RD, Sherbourne CD, Mazel RM. The RAND 36-Item Health Survey 1.0. Health Economics. 1993;2:217-227.

Mosier CI. On the reliability of a weighted composite. *Psychometrika*. 1943;8:161-168.

Nelson EC, Landgraf JM, Hays RD, Wasson JH, Kirk JW. The functional status of patients: how can it be measured in physicians' offices? *Med Care.* 1990;28:1111-1126.

Nunnally J. *Psychometric Theory.* 2nd ed. New York, NY: McGraw-Hill; 1978.

Perrine KR. A new quality-of-life inventory for epilepsy patients: interim results. *Epilepsia*. 1993; 34(suppl 4):S28-S33.

Stewart AL, Sherbourne CD, Hays RD, Wells KB, Nelson EC, Kamberg CJ, Rogers WH, Berry SH, Ware JE. Summary and discussion of MOS measures. In: Stewart AL, Ware JE (eds). Measuring Functioning and Well-Being: The Medical Outcomes Study Approach. Durham, NC: Duke University Press; 1992:345-371.

Vickrey BG, Hays RD, Graber J, Rausch R, Engel J, Brook RH. A health-related quality of life instrument for patients evaluated for epilepsy surgery. *Med Care.* 1992;30:299-319.

Vickrey BG, Perrine KR, Hays RD, Hermann BP, Cramer JA, Meador KJ, Devinsky O. Scoring Manual for the QOLIE-31, Version 1.0. Santa Monica, CA: RAND; 1993.

Ware JE, Sherbourne CD. The MOS 36-Item Short-Form Health Survey (SF-36): I. Conceptual framework and item selection. *Med Care.* 1992; 30:473-483.

ACKNOWLEDGMENTS

Development of QOLIE-89 was supported by an unrestricted research grant from Wallace Laboratories.

Support for the project was provided by Jacki Gordon, PhD, Karen Soderquist, Debra Weiner, and the staff at the Professional Postgraduate Services division of Physicians World Communications Group.

SAS programming and data base management were conducted by Karen Spritzer at RAND.

Sites participating in data collections were (in alphabetical order): Epilepsy Center, University of Alabama School of Medicine, Birmingham, Alabama, Barrow Neurological Institute, Department of Neurology/Epilepsy Center, Phoenix, Arizona, Arizona Comprehensive Epilepsy Program, University of Arizona Health Sciences Center, Tucson, Arizona, Colorado Neurological Institute Epilepsy Center, Englewood, Colorado; Epilepsy Program, Yale University School of Medicine, New Haven. Connecticut: Epilepsy Center, Veterans Affairs Medical Center, West Haven, Connecticut; Comprehensive Epilepsy Center, University of Miami, Miami, Florida; Emory University Clinic, Department of Neurology, Atlanta, Georgia; Department of Neurology, Medical College of Georgia Hospital and Clinics, Augusta, Georgia; Department of Psychology and Social Sciences, Rush-Presbyterian-St. Luke's Medical Center, Chicago, Illinois; Comprehensive Epilepsy Center, Beth Israel Hospital, Boston, Massachusetts; Department of Neurology, New England Medical Center, Boston, Massachusetts; Comprehensive Epilepsy Center, University of Massachusetts Medical Center, Worcester, Massachusetts: Neuropsychology Program and Comprehensive Epilepsy Program, University of Michigan Medical Center, Ann Arbor, Michigan; The Minnesota Epilepsy Group, P.A., St. Paul, Minnesota; Comprehensive Epilepsy Center, New York University Medical Center, Hospital for Joint Diseases. New York, New York; Comprehensive Epilepsy Program, University of Rochester, Rochester, New York; Epilepsy Center, State University of New York at Stony Brook, Stony Brook, New York, Epilepsy Treatment Center, University of Cincinnati Medical Center, Cincinnati, Ohio; Department of Neurology, Section of Epilepsy and Sleep Disorders, The Cleveland Clinic Foundation, Cleveland, Ohio; Oregon Comprehensive Epilepsy Program, Good Samaritan Hospital and Medical Center, Portland, Oregon; Comprehensive Epilepsy Center, Graduate Hospital, Philadelphia, Pennsylvania; Neurosciences Research Institute, Allegheny-Singer Research Institute, Allegheny Campus, Medical College of Pennsylvania, Pittsburgh, Pennsylvania, Epi-Care Center, Baptist Memorial Hospital, Semmes-Murphey Clinic, and University of Tennessee, Memphis, Tennessee, Epilepsy Center, George Washington University Hospital. Washington, D.C.